

The Local Luminosity Function at 25 Microns

David L. Shupe, Fan Fang, Perry B. Hacking

Infrared Processing and Analysis Center, Jet Propulsion Laboratory, Caltech 100-22,
Pasadena, CA 91125

and

John P. Huchra

Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138

Received _____; accepted _____

ABSTRACT

The local luminosity function at $25\ \mu\text{m}$ provides the basis for interpreting the results of deep mid-infrared surveys planned or in progress with space astrophysics missions including ISO, WIRE and SIRTf. We have selected a sample of 1458 galaxies from the IRAS Faint Source Survey with a flux density limit of 250 mJy at $25\ \mu\text{m}$. The local luminosity function is derived using both parametric and non-parametric maximum-likelihood techniques, and the classical $1/V_{\text{max}}$ estimator. Comparison of these results shows that the $1/V_{\text{max}}$ estimate of the luminosity function is significantly affected by the Local Supercluster.

A maximum-likelihood fit to the radial density shows no systematic increase that would be caused by density evolution of the galaxy population. The density fit is used to correct the $1/V_{\text{max}}$ estimate. We also demonstrate the high quality and completeness of our sample by a variety of methods.

The luminosity function derived from this sample is compared to previously published estimates, showing the prior estimates to have been strongly affected by the Local Supercluster. Our new luminosity function leads to lower estimates of mid-infrared backgrounds and number counts.

Subject headings: infrared: sources – luminosity function

1. Introduction

Much of the effort to study infrared-luminous galaxies has centered on wavelengths greater than $50\ \mu\text{m}$. Modeling work is focused on the near-IR (e.g. Chokshi et al. 1994) and far-IR (e.g. Hacking et al. 1987; Rowan-Robinson et al. 1996) portions of the galaxian spectrum. However, the mid-infrared is well-suited for studying starburst and ultraluminous galaxies. About 40% of the luminosity from starburst galaxies is radiated from $8\text{--}40\ \mu\text{m}$ (Soifer et al. 1987). Extinction effects are small, and problems due to infrared cirrus are minimized. Most importantly for space astrophysics, for a fixed telescope aperture, the spatial resolution is higher at shorter wavelengths, and the confusion limit lies at higher redshifts. Recent work using the *Infrared Space Observatory (ISO)* (e.g. Knapp et al. 1996; Boulade et al. 1996; Rowan-Robinson et al. 1996) shows the relative importance of the $7\ \mu\text{m}$ and $15\ \mu\text{m}$ bands for galaxy studies. The *Wide-Field Infrared Explorer (WIRE)*, a Small Explorer mission due to launch in late 1998 (Hacking et al. 1996; Schember et al. 1996), will conduct a very deep survey at $24\ \mu\text{m}$ to study starburst galaxy evolution. The *Space Infrared Telescope Facility (SIRTF)* is also expected to conduct surveys in mid-infrared bands. To interpret the results of these surveys now in progress or soon to commence, it is necessary to better understand the mid-infrared properties of galaxies in the local Universe.

The $25\ \mu\text{m}$ luminosity function provides the basis for predicting the faint source counts in the mid-infrared. The empirical model of Hacking & Soifer (1991) uses an analytic fit to the luminosity function derived by Soifer & Neugebauer (1991). This function was estimated from a complete subsample of the Bright Galaxy Sample (Soifer et al. 1987) containing 135 galaxies to a flux density limit of $1.26\ \text{Jy}$. The availability of many more redshifts of IRAS galaxies (principally from the $1.2\ \text{Jy}$ Survey (Strauss et al. 1990; Fisher et al. 1995)) enables a much larger sample to be studied, reducing uncertainties at high and low luminosities.

In this paper we present the selection of a large galaxy sample that is flux-limited at $25\ \mu\text{m}$, and derive the local luminosity function based on this sample. The sample selection is described in the next section. In Section 3 we describe the $1/V_{\text{max}}$ and the maximum-likelihood estimators for deriving the local luminosity function, and present the results. The completeness of the sample is discussed in Section 4. In Section 5 we calculate the radial density distribution of the sample using a maximum-likelihood method. The radial density fit is used to correct the $1/V_{\text{max}}$ estimate of the local luminosity function, as well as the redshift distribution with which the luminosity function can be compared. Section 6 includes discussions of the different luminosity function estimators, a comparison of our newly derived luminosity function with previous estimates, the implications for mid-infrared backgrounds and number counts, and the effects of evolution on the derivation of the luminosity function. The color properties of the sample are treated in another paper (Fang et al. 1998).

2. Sample Selection

Our sample is based on a selection from the IRAS Faint Source Survey (FSS; Moshir et al. 1992). The principal data product of the FSS is the Faint Source Catalog (FSC). The FSC was produced by coadding IRAS detector scans before extracting sources, and is roughly one magnitude deeper than the Point Source Catalog. Another FSS database is the Faint Source Reject File (FSCR), which contains possible detections that were not included in the FSC for assorted quality-control reasons. This section treats the selection of sources from the FSC and the FSCR in turn.

To minimize the effects of cirrus and contamination from stars, we chose a sky coverage limit of $|b| \geq 30^\circ$. In addition, the same excluded areas used by Strauss et al. (1990) were excluded from our sky coverage region. At high Galactic latitudes, the excluded regions

consist almost wholly of gaps not covered by IRAS, and the Large and Small Magellanic Clouds. Our sample covers 47.7% of the sky.

The 90% completeness limit for the FSC lies at a $25\ \mu\text{m}$ flux density of 210 mJy for sky covered by 2 HCONs, and at 170 mJy for coverage of 3 HCONs, for $|b| > 10^\circ$ (Moshir et al. 1992). Our preliminary searches of redshift databases indicated, however, that for such faint flux density limits, many of the faintest objects would have no redshift available. A flux density limit of 250 mJy was found to have a small redshift incompleteness (see below) and was selected for our sample. Furthermore, sources with moderate (SNR=3 to 5) or high-quality (SNR > 5) flux densities at $25\ \mu\text{m}$ are included in the sample.

A set of weak color criteria were chosen to further discriminate against stars. The great majority of galaxies have larger flux density at $60\ \mu\text{m}$ than at $25\ \mu\text{m}$. For these sources, no constraint was placed on the $12\ \mu\text{m} - 25\ \mu\text{m}$ color. A small percentage of galaxies will have a smaller flux density at $60\ \mu\text{m}$ than at $25\ \mu\text{m}$. For these sources, we have limited the ratio of $F_\nu(25)/F_\nu(60)$ to lie between 1 and 1.6, and have further constrained $F_\nu(12)/F_\nu(25)$ to be less than 1 to avoid admitting large numbers of stars into the sample. Color-color diagrams of the sources in the sample that have been identified as galaxies with redshifts (see below) are displayed in Figure 1. These diagrams show that only a handful of galaxies satisfy $F_\nu(25)/F_\nu(60) \geq 1$, and that our color criteria are fairly liberal for sources identified as galaxies.

To summarize, our sample selection criteria are:

$$\begin{aligned}
 &F_\nu(25) \geq 250\text{mJy}; \\
 &\text{moderate or good quality detection at } 25\mu\text{m}; \\
 &|b| \geq 30^\circ, \text{ and not in Strauss et al. excluded zone;} \\
 &F_\nu(25)/F_\nu(60) < 1 \text{ and no constraint on } F_\nu(12)/F_\nu(25), \\
 &\text{or } 1 < F_\nu(25)/F_\nu(60) < 1.6 \text{ and } F_\nu(12)/F_\nu(25) < 1.
 \end{aligned}$$

1619 sources in the FSC meet the above criteria. To identify galaxies and non-galaxies (stars, infrared cirrus, etc.), and to obtain redshifts for identified galaxies, a positional match of these 1619 sources was made with the 1.2 Jy Survey catalog, the November 1993 public version of J.P. Huchra’s ZCAT, the NED database, and the SIMBAD database. A matching radius of 1 arcminute was used to match the FSC sources with these catalogs. Additional redshifts were culled from J.P. Huchra’s private ZCAT catalog, and from additional observations made at the Fred L. Whipple Observatory (FLWO) 1.5m. This matching process resulted in 1438 galaxies with redshifts. Of the remainder, 166 sources were identified as Galactic objects or HII regions in nearby galaxies, leaving 15 sources identified as galaxies without redshifts.

The selection of sources from the FSCR is more complicated due to the low reliability of sources in that database. Applying our selection criteria to the Faint Source Reject File results in 851 sources. These were matched with the IRAS OPTical IDentification (OPTID) database at IPAC. The OPTID database is a special version of the FSC and FSCR with optical identifications added from the Guide Star Catalog (GSC: Lasker et al. 1990), the Tycho Input Catalog (TIC: Egret et al. 1992)), and the COSMOS/UKST Southern Sky Object Catalogue (COSCAT: Yentis et al. 1992). Sources matching with stars within 1σ error ellipses on the OPTID Plots were excluded. We examined those sources which have no match with either a star or a galaxy within 1σ error ellipses by studying the OPTID Reports and by comparing with the optical Digitized Sky Survey images. We found only a few of these sources to be possible galaxies which, along with the reject sources that match with galaxies within the 1σ error ellipses, gives a small sample of 27 sources. Among these, there are 5 sources which are extremely blue, and have null SNRs at $60\ \mu\text{m}$. ADDSCANS (one-dimensional coadds of IRAS scans) of these sources show no obvious detections at their locations at all four IRAS bands. No redshifts for these sources were found when matching them with the redshift catalogs (see above), and we suspect that they are not real galaxies.

The final yield from the Reject File is 22 sources.

Matching the 22 FSCR sources with the same redshift catalogs gives 20 sources with redshifts. To verify that these matches are not random, a sample of 851 sources randomly distributed in our sampling regions was generated. The number of matches with galaxies is only 2. This indicates that these 22 reject sources are not likely to be just random matches with galaxies. Furthermore, it shows that the number of coincidental matches in our total sample should also be very small.

The breakdown of sources for redshifts for our entire sample (FSC plus FSCR sources) is listed in Table 1. The previously unpublished redshifts obtained at FLWO are tabulated in Table 2. The total number of galaxies with redshifts is 1458. There are 17 galaxies remaining without redshifts.

The 1.2 Jy Survey catalog also provides the ADDSCAN flux densities for the sources extended at 60 μm . For those sources which have no ADDSCAN flux densities in the 1.2 Jy Survey but are extended at 25 μm , we have obtained new ADDSCAN flux density measurements and substituted these for the FSC values.

Figure 2 shows the $\log(N)$ versus $\log(F_\nu(25))$ plot for the sample of galaxies with redshifts. The points follow a slope of -1.5 all the way to our flux limit, demonstrating the high completeness of our sample as a function of flux. The apparent excess at high flux values is most likely due to the Local Supercluster. Further checks of completeness are presented in Section 4, including an investigation of possible systematic effects of flux errors on derived luminosity functions.

For each galaxy, secondary distances are used when available. Otherwise, distances have been assigned using a linear Virgocentric inflow model (Aaronson et al. 1982) and a Hubble constant of 75 $\text{km s}^{-1} \text{Mpc}^{-1}$ (see Appendix A for the case of a different Hubble

constant). The same criteria as Soifer et al. (1987) have been used to identify Virgo cluster galaxies (with a distance of 17.6 Mpc). The monochromatic luminosity of each source is computed from the distance, using a k -correction that assumes a power-law slope of the SED between 12 μm and 25 μm . The monochromatic luminosities are expressed as νL_ν and have units of solar luminosities.

3. Luminosity function results

3.1. $1/V_{max}$ and parametric maximum likelihood

The classic method of estimating luminosity functions is via the $1/V_{max}$ estimator (Schmidt 1968). This technique is non-parametric, so no analytic form of the luminosity function need be assumed, but the data must be binned. The infrared luminosity functions presented by Soifer et al. (1987) and Soifer & Neugebauer (1991) were derived via this method. The space density ρ_L and its error σ_ρ are computed from the following quantities:

$$\rho_L = \left(\frac{4\pi}{\Omega} \right) \left(\sum \frac{1}{V_{max}} \right), \quad (1)$$

$$\sigma_\rho = \left(\frac{4\pi}{\Omega} \right) \left(\sum \frac{1}{V_{max}^2} \right)^2, \quad (2)$$

where Ω is the solid angle of the survey, V_{max} is the maximum volume to which the object could have been detected, and the sum is over all galaxies in each luminosity bin. The assumption that galaxies are distributed uniformly in space may be tested by checking that V/V_{max} is 0.5 in each luminosity bin.

The luminosity function may also be derived via maximum likelihood techniques that are independent of density variations (e.g. Sandage, Tammann, & Yahil 1979). The parametric method has the advantage that no binning of the data is required, but an analytic form of the luminosity function must be assumed. For all of the parametric

maximum likelihood fits in this paper, we have assumed a cumulative luminosity function of the form used by Yahil et al. (1991) in the analysis of the 1.2 Jy Survey:

$$\Psi(L) = C \left(\frac{L}{L_*} \right)^{-\alpha} \left(1 + \frac{L}{L_*} \right)^{-\beta} \quad (3)$$

with its corresponding differential luminosity function

$$\Phi(L) = \left(\frac{\alpha}{L} + \frac{\beta}{L_* + L} \right) \Psi(L). \quad (4)$$

A flux-limited sample contains only a small number of very sub-luminous galaxies, making it difficult to determine the luminosity function at these luminosities. We have followed Yahil et al. (1991) in imposing a lower limit on luminosity $L_s = 4\pi r_s^2 \nu f_m$, where r_s is the distance corresponding to a velocity of 500 km s^{-1} , and f_m is the survey flux limit. For $f_m = 250 \text{ mJy}$, L_s is $4.0 \times 10^7 L_\odot$. Then the minimum detectable luminosity at distance r is $L_{min}(r) = \text{Max}(L_s, 4\pi r^2 \nu f_m)$. The probability of detecting a galaxy of luminosity L_i at distance r_i is then

$$f(L_i|r_i) = \begin{cases} \Phi(L_i)/\Psi(L_{min}) & \text{if } L_i \geq L_{min} \\ 0 & \text{otherwise} \end{cases} \quad (5)$$

The likelihood function is the product of all of these probabilities:

$$\Lambda = -2 \sum_i \ln f(L_i|r_i). \quad (6)$$

This (reciprocal) likelihood function is minimized to determine α , β , and L_* . The variance of these parameters is estimated by making use of the asymptotic normality of the maximum likelihood estimators for a large sample (Kendall, Stuart & Ord 1987).

The likelihood function contains ratios of the differential and cumulative luminosity functions, so another calculation is required to find the normalization C . One method is to use the n_1 estimator of Davis & Huchra (1982) after the shape parameters have been determined. The quantity n_1 is the number density of galaxies with $L \geq L_s$. (See Equations

11 and 13 of Yahil et al. (1991) for the formulas appropriate to our choice of the parametric luminosity function.) An upper limit to the redshift must be chosen to fix the sample volume used in this density estimator. Based on a calculation of n_1 in radial shells, we have chosen an upper limit on redshift of 20,000 km s⁻¹ ($z \sim 0.067$) for the normalization of the maximum likelihood fit. (However, this upper redshift limit is *not* applied in determination of the shape of the luminosity function.) Effects of incompleteness or evolution may become significant beyond this redshift. A maximum-likelihood fit to the density variations of the sample as a function of redshift (see Section 5) further indicates that the density is well-behaved to this limit. A second means of determining the normalization is to require that the number of sources predicted by the luminosity function over some redshift range equals the number in the sample. Computing the normalization from source counts in this way from 500 km s⁻¹ to 20,000 km s⁻¹ gives essentially the same normalization as the n_1 estimator.

The derived parameters for the maximum likelihood fit are listed in Table 3. Luminosity functions derived by the $1/V_{max}$ method and the parametric maximum likelihood method are shown in Figure 3. We show the luminosity functions in the form of the visibility $\Theta = \Phi L^{2.5}$. The visibility is proportional to the number of galaxies visible per log luminosity bin, and it has the advantage of showing the relative numbers of galaxies in a flux-limited sample. The open triangles are from the $1/V_{max}$ estimate of the entire sample, and the solid curve is the parametric maximum-likelihood fit to galaxies with redshifts greater than 500 km s⁻¹, with normalization from a volume-limited sample as described above. The solid curve is truncated at $L_s = 4.0 \times 10^7 L_\odot$. The $1/V_{max}$ estimate (made in bins of $\log(\nu L_\nu) = 0.4$) was converted to visibility by multiplying ρ_L by $L^{1.5} \frac{2.5}{\ln 10}$.

A plot of V/V_{max} as a function of luminosity for our sample is included in Figure 4. The variations in this statistic for the full sample and northern sample at luminosities less

than $10^{10}L_{\odot}$ are most likely due to the Local Supercluster. The southern sample is much more uniform at these luminosities, supporting this interpretation.

The general agreement between the $1/V_{max}$ luminosity function and the shape of the parametric fit in Figure 3, together with the uniformity of the V/V_{max} statistic, confirm that the parametric fit is a good description of the shape of the local luminosity function. The $1/V_{max}$ point at $\log \nu L_{\nu} = 8.6$ corresponds to the luminosity at which the flux limit samples the Virgo cluster, and hence lies well above the solid curve.

The number of galaxies without redshifts (17) is slightly more than one percent of our sample. One method for estimating the effects of this known incompleteness on the luminosity function is to assign to each of these galaxies the median redshift of their log flux bin, and recalculate the luminosity function. When the median redshifts are computed for the whole sample, all the assigned redshifts are around 7,000 km sec⁻¹. When the median redshifts are calculated from those galaxies whose redshifts were not obtained from public sources (a more realistic estimate), the assigned redshifts range from 12,000 to 18,000 km sec⁻¹. Since all these galaxies have a 25 μ m flux density within a factor 1.5 of the flux limit, their luminosities all fall near L_* . When the luminosity function is recalculated and plotted as a visibility, we have found that the asymptotes are unchanged, and the peak visibility is increased by a few percent.

3.2. Nonparametric maximum likelihood

A non-parametric maximum-likelihood method (a.k.a “Stepwise Maximum Likelihood”) has been used to calculate the galaxy luminosity function in various surveys (Lynden-Bell 1971; Choloniewski 1986; Efstathiou et al. 1988; Saunders et al. 1990; Marzke et al. 1994). This method does not assume a specific functional form for the luminosity function, but

rather describes it by a non-negative step function (i.e. requires binning) which makes the observed galaxy luminosity distribution as likely as possible. We have followed the prescriptions set out by Efstathiou et al. (1988) and Saunders et al. (1990) for calculating the non-parametric luminosity function.

As in the parametric maximum-likelihood method, the non-parametric approach also eliminates the effect of any density inhomogeneities, but loses the overall normalization by assuming that the galaxy spatial and luminosity distributions are independent. The normalization of the non-parametric function has been chosen to give the same source-counts as the parametric function over the same bins.

Figure 5 shows the visibility function obtained by using the non-parametric maximum-likelihood method (circles), which agrees within the error bars with our parametric results (solid line, from Figure 3). The non-parametric method is commonly used to assess whether the parametric functional form is acceptable. Figure 5 shows that the parametric form of Figure 3 gives a reasonable description of the density-independent $25\ \mu\text{m}$ luminosity function. We have found that the non-parametric function is sensitive to the bin width, however. Figure 5 shows the case for bins in $\log L = 0.2$, but for a bin width of 0.4, the non-parametric results is systematically higher than the parametric curve for high luminosities. The coarser binning probably results in a loss of information and a less reliable luminosity function. The choice of bin size is a compromise between this effect and tolerating the statistical uncertainty from having too few sources in each bin.

4. Completeness Checks

Systematic variations of noise across the sky may change the detection quality and cause incompleteness problems for the sample. In the $25\ \mu\text{m}$ passband of IRAS, noise

is dominated by photon shot noise from the zodiacal background, and hence is a strong function of the ecliptic latitude. We have examined the noise variation in the sample and in the corresponding FSS plates and found two regions that contain higher noise at $25\mu\text{m}$. They are essentially the two quadrants defined by $(0^\circ < l < 180^\circ, b < -30^\circ)$ and $(180^\circ < l < 360^\circ, b > 30^\circ)$, which contain the ecliptic plane. Most of the moderate-quality detection sources (generally, those with SNRs between 3 and 5) in our sample are in these two regions, indicating the systematic effect.

To estimate this systematic effect on our sample completeness, we compared luminosity functions estimated for two subsamples of our 250-mJy-limited full sample. The first subsample is defined by raising the flux-density limit to 400 mJy in the two quadrants of the sky $(0^\circ < l < 180^\circ, b < -30^\circ)$ and $(180^\circ < l < 360^\circ, b > 30^\circ)$ where the noise is systematically higher, and keeping the 250 mJy limit in the other two quadrants. Most of the moderate quality detection sources are left out of this subsample, so we call these 1050 sources the high-quality subsample. The second subsample consists of those 408 sources in the full sample that are left out of the high-quality subsample, and is therefore a low-quality subsample. Figure 6 compares the luminosity functions of the high-quality and the low-quality subsamples, using the $1/V_{max}$ estimator since it accomodates the different flux limits more easily than either of the maximum-likelihood estimators. There is no systematic deficiency of galaxies in the low-quality subsample. The luminosity functions agree within the 1σ error bars, with the exception of the point at $\log \nu L_\nu = 8.6$. The low-quality subsample includes galaxies in the Virgo cluster with $25\mu\text{m}$ flux densities between 250 and 400 mJy, most of which fall into this luminosity bin.

In Figure 7 we compare the $1/V_{max}$ luminosity functions of the high-quality subsample and the full 250 mJy-limited sample. They agree very well, indicating that the full sample is complete at 250 mJy flux-density across the entire sampling sky. Therefore we are confident

that the full 250 mJy-limited sample is well-defined.

As another check of the completeness of our full sample, we have computed the cumulative probability of finding a galaxy of luminosity L at distance r , following the method of Yahil et al. (1991). This probability is the integral of Equation 5 and is $F(L|r) = \Psi(L)/\Psi(L_{min}(r))$. This probability should be uniformly distributed on the interval $[0,1]$ independent of distance. If the sample were incomplete at the faint end, we would expect a decrease in the relative number of galaxies as $F \rightarrow 1$.

We have computed $F(L_i|r_i)$ for each galaxy in the sample, then computed the histogram of this statistic in different distance ranges, normalized to 1 in each range. The result is shown in Figure 8. There is no systematic decrease in the relative galaxy counts as $F \rightarrow 1$ (albeit the statistics are poor beyond 400 Mpc). This result provides further confirmation that our 250 mJy-limited sample is complete to these distances. Since this method relies upon ratios of the luminosity function, it should be insensitive to inhomogeneities in the galaxy distribution.

5. Density Variations and the Corrections for V/V_{max}

Just as the galaxy luminosity function independent of the inhomogeneities of the density field can be obtained, the spatial density variations independent of the galaxy luminosity function can also be calculated using the maximum-likelihood method (Saunders et al. 1990). Here we only investigate the radial density distribution. The probability for detecting a source of luminosity L_i at radial distance r_i is

$$p_i = \frac{\rho(r_i)dV_i}{\int_0^{r_{max,i}} \rho(r)(dV/dr)dr}, \quad (7)$$

where $\rho(r_i)$ is the density at r_i , and $r_{max,i}$ is the maximum distance for the source to be included in the sample. The likelihood function for all the detected sources would

be $\mathcal{L} = \prod p_i$, given the density distribution $\rho(r)$ we are trying to calculate. In the non-parametric approach, this distribution can be found by maximizing $\ln \mathcal{L}$ with respect to the binned density field $\rho(r_j)$, which gives

$$\rho(r_j) = N_j \left(\sum_i \frac{f_{ij} V_j}{V_{eff,i}} \right)^{-1}, \quad (8)$$

where $V_{eff,i} = \sum_{k=0}^{r_{max,i}} \rho(r_k) V_k$, N_j is the number of sources in the j -th distance bin with volume V_j , f_{ij} is the fraction of the bin contained in $[0, r_{max,i}]$ where the i -th source is observable, and the sum is over all sources.

Figure 9 illustrates the non-parametric maximum-likelihood estimate of the radial density variation (solid line) as compared with the flux-limited observed distribution (dashed line). Here the radial distance bins are in logarithmic scale, and the density variations are normalized to the average density over all bins of interest. The method correctly recovers the density enhancement at ~ 17 Mpc caused by Virgo.

We do not find a systematic increase in the radial density at large distances. If our sample is assumed to be complete, this result means that there is no indication of density evolution in this sample. We return to this point in section 6.4.

The benefits of calculating the radial density variations in our sample are two-fold. First, it may be used to correct the V/V_{max} statistic and the traditional $1/V_{max}$ estimate of the density-dependent luminosity function (Schmidt 1968; Felton 1977). For the V/V_{max} test, since

$$\frac{\int_0^{r_{max}} \frac{\int_0^r \rho(r') D^2 dr'}{\int_0^{r_{max}} \rho(r') D^2 dr'} \rho(r) D^2 dr}{\int_0^{r_{max}} \rho(r) D^2 dr} = \frac{\int_0^{r_{max}} (\int_0^r \rho(r') D^2 dr') d(\int_0^r \rho(r) D^2 dr)}{(\int_0^{r_{max}} \rho(r) D^2 dr)^2} = \frac{1}{2}, \quad (9)$$

where $\rho(r)$ is the radial density field, r_{max} is the maximum comoving distance for a source to be included in a sample, and D is the effective distance (Longair 1978), we can modify the V/V_{max} test so that

$$\frac{V}{V_{max}} = \frac{\int_0^r \rho(r) D^2 dr}{\int_0^{r_{max}} \rho(r) D^2 dr}, \quad (10)$$

which gives $\overline{V/V_{max}} = 1/2$ for an inhomogeneous radial distribution. This also suggests that we use the $V_{eff,i}$ in equation 8 instead of V_{max} in the $1/V_{max}$ estimator to calculate a density-corrected luminosity function. The $\rho(r)$ calculated from the maximum-likelihood method can provide these corrections. The results are shown in Figure 10 for the new V/V_{max} values and in Figure 11 for the $1/V_{max}$ luminosity function. Compared with the results without the inhomogeneity correction in Figure 4, the V/V_{max} values are closer to 0.5 for those bins most strongly affected by the Local Supercluster. (However, the corrected V/V_{max} values are not suitable as a test of the sample completeness, since the density correction will mask any incompleteness in the sample.) The new $1/V_{max}$ luminosity function is smoother and closer to the ones estimated by the maximum-likelihood method. As mentioned in Saunders et al. (1990), the binning of the distance in Figure 9 may account for the residual effects of the density inhomogeneities in these Figures.

Second, the “true” density variations in Figure 9 also correct the observed redshift distribution. The density-variation-corrected redshift distribution should be predicted by a density-independent luminosity function estimated in the same sample. This provides an independent test to verify the luminosity function. Figure 12 shows our results. Here the dashed histogram is the observed redshift distribution. (Here we show the source-counts in distance bins instead of redshift bins since some galaxies have secondary distances.) It is divided by the density of the corresponding distance bins in Figure 9 which gives the solid-line histogram. The source-counts affected by large-scale structures, such as the ones near Virgo, are corrected by this procedure. The dotted line shows the prediction by the parametric density-independent luminosity function of Figure 3 and Table 3. This prediction fits the solid-line histogram well, demonstrating the reliability of our luminosity function. Both the solid-line histogram and the dotted line are normalized so that they contain the same total number of sources as observed within $20,000 \text{ km s}^{-1}$.

6. Discussion

6.1. Limitations of the methods

The preceding sections have presented the 25 μm luminosity function calculated by a number of methods. In this section, the limitations of the various methods are revisited in greater detail.

The $1/V_{max}$ estimate of the luminosity function varies the most from the other estimators, particularly at luminosities less than L_* , where our sample is most strongly influenced by the Local Supercluster. As noted above, this estimator assumes that the 3-dimensional galaxy distribution is homogeneous. The error bars in all of our $1/V_{max}$ estimates similarly depend on the assumption of uniformity, and do not reflect the additional uncertainties that arise from an inhomogeneous distribution. We consider the $1/V_{max}$ estimates to be the least reliable descriptions of the 25 μm luminosity function out of those we have presented.

The parametric and non-parametric maximum-likelihood estimates should be independent of structures in the galaxy distribution. However, these estimators assume that the luminosity function is independent of density. This assumption is difficult to check. In the case of a large 60 μm -selected sample, Yahil et al. (1991) demonstrated that their similarly-derived luminosity function provided a good description of the luminosity distribution of galaxies even in regions of high density.

The non-parametric maximum likelihood method applied to our sample is sensitive to the choice of binning, as described in section 3.2. We therefore consider the parametric maximum-likelihood description of the 25 μm luminosity function with parameters in Table 3 to be the best out of those we present in this paper.

6.2. Comparison with other luminosity functions

As mentioned in the Introduction, Soifer & Neugebauer (1991) derived the local luminosity function at $25\ \mu\text{m}$ from a complete subsample of the $60\text{-}\mu\text{m}$ -selected Bright Galaxy Sample (Soifer et al. 1987). This subsample contained 135 galaxies to a flux density limit of 1.26 Jy, and the luminosity function was determined using the classical $1/V_{\text{max}}$ estimator. Their luminosity function is plotted as a space density in Figure 13, together with our parametric and $1/V_{\text{max}}$ estimates.

Our flux density limit of 250 mJy samples the Virgo cluster at $\log \nu L_\nu = 8.6$ (section 3.1), and our $1/V_{\text{max}}$ estimate is much higher than the parametric curve in that bin. The Soifer & Neugebauer points in Figure 3 in $\log \nu L_\nu = 9.4$ bin is also much higher than the curve, as expected since their flux density limit is five times higher than ours. We conclude that the Soifer & Neugebauer luminosity function is strongly affected by the Local Supercluster, particularly because the Bright Galaxy Sample was selected to be visible from Palomar Observatory (Soifer et al. 1984) and does not include the south Galactic cap.

6.3. Implications

The improved luminosity function that is the main result of this paper has implications the luminosity density of the local Universe, infrared number counts predictions at $25\ \mu\text{m}$, and the infrared background expected at this wavelength.

Soifer & Neugebauer (1991) calculated the local luminosity density in the four IRAS passbands based on their derived luminosity functions. At $25\ \mu\text{m}$, they found that the log of the luminosity density in units of $L_\odot\ \text{Mpc}^{-3}$ is 7.2. Integrating our parametric function gives a log luminosity density of 6.9, which is a factor of 2 lower in linear terms than the Soifer & Neugebauer result. The differences in the sub- L_* portion of the luminosity

functions account for the different integrated luminosity densities.

Hacking & Soifer (1991) presented number count distributions at $25\ \mu\text{m}$, using a fit to the Soifer & Neugebauer (1991) luminosity function. (Those authors fit a hyperbola to the visibility function, which has the power-law asymptotes at high and low luminosities like our parametric form, except that it forces the absolute values of the slopes of the asymptotes to be identical.) As they noted, the Soifer & Neugebauer functions at other wavelengths are 20%-30% higher than those derived from other samples (Saunders et al. 1990). Our improved $25\ \mu\text{m}$ function is similarly about 15% lower in normalization than the Soifer & Neugebauer estimate. Hence, models based on this luminosity function and its normalization will produce number counts about 15% lower at the bright end than those in Figure 1(c) of Hacking & Soifer (1991). Such a change would bring the number counts in better agreement with the IRAS number counts presented in their same figure.

Finally, we have substituted our luminosity function into the backgrounds model of Hacking & Soifer (1991). The modified model yields estimates of the infrared background at $25\ \mu\text{m}$ that are factors of 2.5 to 2.7 times smaller than the original Hacking & Soifer estimates. Our results indicate that the detection of an IR background at this wavelength is likely to be even more difficult than previously predicted.

6.4. Evolution

To this point, we have neglected any effects of galaxy evolution on our derivation of the luminosity function. Since the maximum-likelihood estimators are independent of density variations, the shape of the luminosity function will be unchanged by density evolution of the galaxy population. Our normalization considers galaxies out to $20,000\ \text{km s}^{-1}$, a redshift where evolution effects should be small.

More importantly, our maximum likelihood fit to the radial density (section 5 and Figure 9) does not show a systematic increase with redshift and hence no indication of density evolution. This result is in direct contrast to that of Saunders et al. (1990). From their 60- μm -selected sample, those authors found that density evolution of $(1+z)^{6.7}$ matched their radial density fit. However, we note that the existence of evolution from IRAS samples is not yet settled — for the deepest IRAS sample, Ashby et al. (1996) did not find the high-redshift tail expected from strong evolution.

Nevertheless, we have computed the shape of the parametric luminosity function for the case of exponential luminosity evolution corresponding to $(1+z)^3$ at low redshifts (cf. Saunders et al. (1990)), by correcting the luminosity of each galaxy to the present epoch. The values of the shape parameters of this function are $\alpha = 0.437 \pm 0.032$, $\beta = 1.769 \pm 0.067$, and $L_* = 3.98 \pm 0.34 \times 10^9 L_\odot$. The only change from the function described in Table 3 is that the slope of the luminosity function at the high-luminosity end is slightly lowered.

7. Summary and conclusions

The following are the results of this paper:

1. We have selected a sample of 1458 galaxies with redshifts from the IRAS Faint Source Survey with a flux density limit of 250 mJy at 25 μm . An additional 17 galaxies do not have redshifts available.
2. The local luminosity function is derived using the $1/V_{max}$ estimator and both parametric and non-parametric maximum likelihood methods. The $1/V_{max}$ estimate is significantly affected by the Local Supercluster. The maximum likelihood methods are independent of density variations, and we consider the parametric fit with parameters in Table 3 to be the best estimate of the local luminosity function at 25 μm .

3. A maximum likelihood fit to the radial density in this sample is used to correct the $1/V_{max}$ estimate. The fit shows no sign of a systematic increase with redshift of the density, as would result from density evolution of the galaxy population.

4. The $1/V_{max}$ luminosity function derived from a smaller sample by Soifer & Neugebauer (1991) is significantly contaminated by the Local Supercluster. Predictions of number counts and local luminosity density based on that function are 15-20% higher than those indicated by our improved luminosity function. The new function also leads to lower predictions of the mid-infrared background due to galaxies.

We thank Susan Tokarz, Perry Berlind, and Jim Peters for observations made at F.L. Whipple Observatory. We also thank Iffat Khan and Joe Mazzarella for assistance with database searches, and Carol Lonsdale and Tom Soifer for helpful discussions. We are grateful to Cong Xu for a critical reading of this manuscript. We thank the anonymous referee and our Scientific Editor, Steven Willner, for numerous comments that have improved this paper. The SIMBAD database is maintained by CDS in Strasbourg, France. The NED database is supported at IPAC by NASA.

A. The Distance to Virgo and a Different Hubble Constant

We have assumed a distance 17.6 Mpc to Virgo in the linear Virgocentric inflow model (Aaronson et al. 1982) to calculate the distance to each galaxy which does not have a secondary distance. To be consistent with our assumption of $75 \text{ km s}^{-1} \text{ Mpc}^{-1}$ for the Hubble constant, we have used 1019 km s^{-1} as the observed velocity of Virgo corrected to the centroid of the Local Group, and implicitly assumed an infall velocity of about 300 km s^{-1} for the Local Group toward Virgo. The resulting Hubble flow velocity of Virgo, $\sim 1320 \text{ km s}^{-1}$, is consistent with many other estimates (e.g. Faber et al. 1989; Tonry et

al. 1990; Mould et al. 1995; Huchra 1996). A different Hubble constant would change the distance estimate to Virgo if the Hubble flow velocity of Virgo remains unchanged.

We have calculated the $1/V_{max}$ luminosity function assuming no Virgocentric inflows, in which we used the redshift as the direct distance measure unless a secondary distance exists for a galaxy. Figure 14 compares the $1/V_{max}$ luminosity functions for the 250 mJy-limited sample with and without Virgocentric inflow. A Hubble constant of $75 \text{ km s}^{-1} \text{ Mpc}^{-1}$ is used in both cases. The two estimates agree very well. This indicates that the inflow model we use has a small impact on the estimate of the distance of the galaxies in the sample, and that most of these galaxies are not strongly affected by the gravity from Virgo. Note that a luminosity function simply scales with the Hubble constant (the luminosity changes as H_0^{-2}) if all distance estimates in a sample are from redshifts.

The number of galaxies with secondary distances in our sample is small (only 22). We have done experiments to verify that these galaxies do not contribute statistically significantly to the calculated luminosity function. We therefore conclude that our results can be simply scaled if a different Hubble constant is used.

REFERENCES

- Ashby, M.L.N., Hacking, P.B., Houck, J.R., Soifer, B.T., & Weisstein, E.W. 1996, ApJ, 465, 428.
- Aaronson, M., Huchra, J., Mould, J., Schechter, P. L., and Tully, R. B. 1982, ApJ, 258, 64.
- Boulade, O. et al. 1996, A&A, 315, L85.
- Chokshi, A., Lonsdale, C. J., Mazzei, P., and De Zotti 1994, ApJ, 424, 578.
- Choloniewski, J. 1986, MNRAS, 223, 1.
- Davis, M., & Huchra, J. 1982, ApJ, 254, 437.
- Efstathiou, G., Ellis, R. S., & Peterson, B.A. 1988, MNRAS, 232, 431.
- Egret, D., Didelon, P., McLean, B. J., Russell, J. L., and Turon, C. 1992, A&A, 258, 217
- Faber, S. M., Wegner, G., Burstein, D., Davies, R. L., Dressler, A., Lynden-Bell, D., and Terlevich, R. J. 1989, ApJS, 69, 763.
- Fang, F., Shupe, D.L., Xu, C., and Hacking, P.B. 1998, ApJ, submitted.
- Fairall, A., Lowe, L. & Dobie, P. 1988, Pub. of the Department of Astronomy of the University of Cape Town, No. 5, *A Catalog of Galaxies South of Declination -30 That Have Been Observed Spectroscopically*.
- Felton, J. E. 1977, AJ, 82, 861.
- Fisher, K. B., Huchra, J.P, Strauss, M., Davis, M., Yahil, A., & Schlegel, D. 1995, ApJS, 100, 69.
- Hacking, P. B. et al. 1996, The Wide-Field Infrared Explorer (WIRE) Mission, to appear in International Symposium on Diffuse Infrared Radiation and the IRTS, Institute of Space and Astronomical Sciences, Sagamihara, Japan.
- Hacking, P. B., Condon, J. J., and Houck, J. R. 1987, ApJ, 316, L15.

- Hacking, P. B., and Soifer, B. T. 1991, *ApJ*, 367, L49.
- Huchra, J., Davis, M., Tonry, J. & Latham, D. 1983, *ApJS*, 52, 89.
- Huchra, J.P. 1996, in *IAU Symposium No. 168, Galaxy Motions in the Nearby Universe*, pp. 143.
- Kendall, M.G., Stuart, A., & Ord, J.K. 1987, *The Advanced Theory of Statistics*, Vol. 2 (5th ed.; New York; Oxford University Press).
- Knapp, G. R., Rupen, M. P., Fich, M., Harper, D. A., Wynn-Williams, C. G. 1996, *A&A*, 315, L75.
- Lasker, B. M., Sturch, C. R., McLean, B. J., Russell, J. L., Jenkner, H., & Shara, M. M. 1990, *ApJ*, 99, 2019
- Lawrence, A., Rowan-Robinson, M., Crawford, J., Parry, I., Xia, X-Y., Ellis, R.S., Frenck, C.S., Saunders, W., Efstathiou, G. and Kaiser, N., 1994 *MNRAS* to be submitted;preprint
- Longair, M. S. 1978, *Observational Cosmology*, A. Maeder, L. Martinet, and G. Tammann (Geneva Observatory).
- Lynden-Bell, D. 1971, *MNRAS*, 155, 95.
- Marzke, R.O., Geller, M.J., Huchra, J.P., & Corwin, H.G. 1994, *AJ*, 108, 437.
- Moshir M., et al. 1992, *IRAS Faint Source Catalog Explanatory Supplement*, Version 2.
- Mould, J. et al. 1995, *ApJ*, 449, 413.
- Rowan-Robinson, M. et al. 1996, *Spectral Energy Distributions, Starburst Models and Star Formation History*, *MNRAS*, submitted.
- Sandage, A., Tamman, G. A., & Yahil, A. 1979, *ApJ*, 232, 352.
- Saunders, W., Rowan-Robinson, M., Lawrence, A., Efstathiou, G., Kaiser, N., Ellis, R. S., & Frenk, C. S. 1990, *MNRAS*, 242, 318.

- Schember, H., Kemp, J., Ames, H., Hacking, P., Herter, T., Fafaul, B. Everett, D., Sparr, L. 1996, “The Wide-Field Infrared Explorer (WIRE)”, SPIE Proceedings, 2744, Infrared Technology and Applications XXII, p. 751-760.
- Schmidt, M. 1968, ApJ, 151, 393.
- Soifer, B. T. et al. 1987, ApJ, 320, 238.
- Soifer, B. T., & Neugebauer, G., 1991, AJ, 101, 354.
- Strauss, M., Davis, M., Yahil, A., & Huchra, J. P. 1990, ApJ, 361, 49.
- Tonry, J., Ajhar, E., & Luppino, G. 1990, AJ, 100, 1416.
- Yahil, A., Strauss, M. A., Davis, M., & Huchra, J. P. 1991, ApJ, 372, 380.
- Yentis et al. 1992, in “Digitised Optical Sky Surveys”, p.67, ed. MacGillivray & Thomson (Dordrecht:Kluwer).

Fig. 1.— Color-color plots for the galaxies with redshifts in the sample. The upper plot shows galaxies with good or moderate detections in each of the 12, 25 and 60 μm bands. The lower plot show galaxies with upper limits in either or both the 12 or 60 μm bands. The hatched region shows the region of the color-color diagram that is excluded from the sample selection.

Fig. 2.— Differential number counts versus flux for the galaxies with redshifts in the sample. The line has a slope of -1.5.

Fig. 3.— Visibility plots for the 250 mJy sample. The open triangles represent the results of the $1/V_{\text{max}}$ estimate of the luminosity function fit to the entire sample. The solid line is the parametric maximum likelihood estimate for galaxies with redshifts greater than 500 km s^{-1} . The parameters of the luminosity function are listed in Table 3. The curve is truncated at $L_s = 4 \times 10^7 L_\odot$. The distance scale at top is computed from the 250 mJy flux density limit and the luminosity scale at the bottom.

Fig. 4.— V/V_{max} for the full sample (upper panel), and separately for northern and southern Galactic latitudes (lower panel). The dashed line at $V/V_{\text{max}} = 0.5$ shows the expected value for a uniform distribution of galaxies.

Fig. 5.— The non-parametric maximum-likelihood estimate of the visibility function (circles). The parametric maximum-likelihood estimate from Figure 3 is also shown for comparison. The different estimators agree within the 1σ error bars shown on the Figure.

Fig. 6.— The $1/V_{\text{max}}$ visibility functions for the high-quality (open squares) and low-quality subsamples (filled squares) defined in Section 4. The 1σ error bars are shown.

Fig. 7.— The $1/V_{\text{max}}$ visibility functions for the high-quality subsample (filled-squares) defined in Section 4 and the full 250 mJy-limited sample (filled-circles). Also shown are the

1σ error bars.

Fig. 8.— Distribution of the quantity $\Psi(L)/\Psi(L_{min})$ in the sample for galaxies in different distance ranges. The mean value is normalized to 1 in each distance range. The flatness of these distributions confirms that the completeness of the sample is high to at least 400 Mpc.

Fig. 9.— The non-parametric maximum-likelihood estimate (solid histogram) of the radial density distribution for the 250 mJy flux-limited sample at $25\ \mu\text{m}$. This estimate is independent of the galaxy luminosity function. The dashed lines show the observed radial density distribution.

Fig. 10.— The V/V_{max} plot for the full flux-limited sample after correcting for density variations as described in the text. The 1σ error bars are shown.

Fig. 11.— The density-corrected $1/V_{max}$ estimate of the visibility function. The parametric maximum-likelihood estimate from Figure 3 is also shown for comparison. The density inhomogeneities have been corrected according to the radial density estimate shown in Figure 9. The different estimators agree within the 1σ error bars shown on the Figure.

Fig. 12.— The observed redshift distribution (dashed-line histogram) and the distribution corrected for the density variations in Figure 9 (solid-line histogram) as described in the text. The dotted line is the distribution predicted by the parametric maximum-likelihood luminosity function from Figure 3 and Table 3.

Fig. 13.— Local luminosity functions at $25\ \mu\text{m}$ plotted in terms of space density as a function of luminosity. The dashed lines were computed by varying the values in Table 3 by their 1σ uncertainties.

Fig. 14.— The $1/V_{max}$ visibility functions for the 250 mJy-limited sample with (filled circles,

as in Figure 7) and without (circles) the Virgo-centric inflow. The two functions agree well within the 1σ error bars, indicating that most of the galaxies in the sample are not affected by the inflow model.

Table 1. Source of redshifts for sample

Source of redshift	Number
Public Sources:	
1.2 Jy catalog	1277
Nov. 1993 ZCAT	49
NED	79
SIMBAD	6
Fairall, Lowe, & Dobie (1988)	1
Private Sources:	
FLWO 1.5m (Table 2)	41
QDOT (Lawrence et al. 1994)	5

Table 2. Previously unpublished redshifts from FLWO

R.A.	Dec.	$F_\nu(25)$	cz
(B1950)		(Jy)	(km s ⁻¹)
00 17 02.7	−04 56 46	0.3734	6181 ± 28
00 20 47.6	+10 29 50	0.2633	69094 ± 85
00 37 17.8	+08 29 18	0.2619	17482 ± 34
00 44 30.0	−18 03 33	0.2776	43979 ± 48
00 44 37.2	+10 14 46	0.3265	$49959 \pm 54(N)$ $50133 \pm 43(S)$
00 56 04.2	+08 31 54	0.2861	17269 ± 36
01 27 44.3	+07 53 04	0.2541	33899 ± 44
01 47 43.0	−16 55 12	0.2875	48432 ± 39
07 55 56.0	+50 58 36	0.3589	16305 ± 41
08 11 42.0	+46 13 00	0.2951	12277 ± 136
08 46 21.1	+11 26 03	0.2626	23218 ± 33
08 57 06.0	+18 01 00	0.2645	$15940 \pm 20(E)$ $15830 \pm 19(W)$
09 39 16.1	+23 54 38	0.3299	6407 ± 39
09 42 35.4	+17 51 45	0.4363	38440 ± 41
10 03 12.0	+13 12 00	0.4967	2789 ± 28
10 07 43.0	+14 27 39	0.2729	59689 ± 55
10 08 05.4	+06 26 45	0.2883	29445 ± 39
10 23 48.0	+00 56 00	0.3076	6422 ± 18

Table 2—Continued

R.A.	Dec.	$F_\nu(25)$	cz
(B1950)		(Jy)	(km s ⁻¹)
11 06 26.0	−23 21 57	0.3648	24551 ± 41
11 19 58.7	+04 31 06	0.5010	11399 ± 38
11 22 24.9	+02 56 02	0.2533	18813 ± 17(<i>N</i>) 18857 ± 24(<i>S</i>)
11 31 56.3	+22 47 49	0.3011	9230 ± 34
11 57 29.2	−03 30 20	0.2663	8024 ± 20
12 12 17.0	−03 12 00	0.4950	10108 ± 37
12 17 18.6	+09 12 59	0.2743	7398 ± 31
12 19 41.7	+00 09 03	0.2991	51652 ± 29(<i>N</i>) 51626 ± 36(<i>S</i>)
12 23 25.0	−23 19 36	0.3232	14568 ± 40
12 36 16.3	−27 02 00	0.2678	7502 ± 22
12 38 18.7	+27 50 19	0.3110	16947 ± 34
12 50 30.4	−02 56 36	0.4896	6807 ± 34
13 24 15.4	−06 08 38	0.2963	13489 ± 46
13 53 28.3	−07 51 36	0.3494	22750 ± 40
13 55 57.1	−15 53 51	0.3377	10868 ± 34
15 19 54.4	−13 51 29	0.2555	7182 ± 25
15 21 47.0	−11 02 07	0.2667	15447 ± 37
15 21 48.0	+05 01 00	0.2588	10715 ± 10

Table 2—Continued

R.A.	Dec.	$F_\nu(25)$	cz
(B1950)		(Jy)	(km s ⁻¹)
21 11 45.0	−27 31 29	0.2655	25801 ± 33
22 34 59.5	−25 44 16	0.2646	23353 ± 21
23 13 02.6	−00 58 50	0.3414	8318 ± 36
23 29 00.6	−12 52 29	0.3778	6357 ± 40
			11478 ± 23(<i>W</i>)
23 40 45.2	+04 16 37	0.3489	11564 ± 21(<i>N</i>)
			11334 ± 21(<i>S</i>)

Note. — Multiple redshift components are labeled with letters denoting their relative orientations.

Table 3. Fitted Parameters of Luminosity Function

α	β	L_* ($L_\odot$)	C (10^{-3} Mpc^{-3})	n_1 (10^{-3} Mpc^{-3})
0.437 ± 0.032	1.749 ± 0.067	$4.07 \pm 0.36 \times 10^9$	1.23 ± 0.09	9.2 ± 0.7



























